

M/G/ ∞ WITH BATCH ARRIVALS

J. KEILSON

GTE Laboratories and Massachusetts Institute of Technology, Cambridge, MA, USA

A. SEIDMANN

University of Rochester, Rochester, NY, USA

Received September 1987

Revised June 1988

Let $p_\infty(n)$ be the distribution of the number $N(\infty)$ in the system at ergodicity for systems with an infinite number of servers, batch arrivals with general batch size distribution and general holding times. This distribution is of importance to a variety of studies in congestion theory, inventory theory and storage systems. To obtain this distribution, a more general problem is addressed. In this problem, each epoch of a Poisson process gives rise to an independent stochastic function on the lattice of integers, which may be viewed as stochastic impulse response. A continuum analogue to the lattice process is also provided.

queues • M/G/ ∞ • batch arrivals

Introduction

The distribution of the number in the system at ergodicity for systems with an infinite number of servers, batch arrivals with general batch size distribution and general holding times is of importance to a variety of studies in congestion theory, inventory theory and storage systems. To obtain this distribution, a more general problem is addressed. In this problem, each epoch of a Poisson process gives rise to an independent stochastic response function on the lattice of integers.

The simplicity and importance of the results obtained suggest that they may be known, but the authors have not been able to find them in the literature. Interesting transform results in a very general setting have been given by Shanbhag [5] and time dependent distributions have been provided by Brown and Ross [1] for the number of customers in the system. Neither paper, however, provides an explicit form for the ergodic distribution of interest. For related results, the reader is referred to the book [2] by Chaudry and Templeton. The method employed in this note is probabilistic and succinct and may be of independent interest.

1. A more general problem

The work described has been motivated by M/G/ ∞ batch arrival system needs. The results however relate simply to a more general system context and may have value elsewhere.

For the M/G/ ∞ batch system, customers arrive at Poisson epochs of rate λ with random batch size and random i.i.d service times to a system having an infinite number of servers. The p.g.f. at ergodicity of $N(t)$, the number in the system, is wanted. To find this distribution, we consider the more general problem where, for each arrival epoch τ_k of the Poisson stream, there is a stochastic response function $N_k^*(y)$ having non-negative integer values with $N_k^*(y) \rightarrow 0$ as $y \rightarrow \infty$. This means that, for a given realization ω , $n_k^*(\omega, y)$ is the number of customers in the batch arriving at τ_k still in the system at time $\tau_k + y$. The random response functions $N_k^*(y)$ are independent and identically distributed in law. The number in the system $N(t)$ is then given by

$$N(t) = \sum_{k=-\infty}^{k=+\infty} N_k^*(t - \tau_k).$$

To obtain the p.g.f. of $N(\infty)$ for the general

problem, one notes that because of the Poisson character of the arrival process, one may regard the input stream as a superposition of M independent sub-streams. Each sub-stream has Poisson arrivals of rate λ/M and has the same stochastic response function $N^*(y)$ and each sub-stream may be thought of as associated with its own subsystem. The response function $N^*(y)$ will be assumed initially to terminate at the level 0 after a service interval of finite duration D . Let $r_n(y) = P\{N^*(y) = n\}$ be the probability that n customers of the batch remain in the system at time y after the arrival of the batch. During the service interval of length D , the batch response function $N^*(y)$ will then be described by the p.g.f.

$$g(u, y) = \sum_n r_n(y) u^n. \tag{1}$$

Because the arrival process for each subsystem is Poisson, the length of the intervals between the termination of service on the previous batch and the start of service on the next batch is exponentially distributed with mean M/λ . Let $I_n(y)$, $n \geq 0$, be the indicator function for the number in a batch remaining at time y being n , i.e. let $I_n(y) = 1$ when $N^*(y) = n$ with $I_n(y) = 0$, otherwise. For a Poisson variate K of parameter θ , $P\{K \geq 2\} = O(\theta^2)$, $\theta \rightarrow 0$. The number of batches present in a substream is Poisson with parameter $(\lambda/M)E\{T_B\}$ where T_B is the batch persistence time. Hence the probability that two or more batches are present simultaneously in a subsystem is $O(\lambda^2/M^2)$ as $M \rightarrow \infty$. For any substream, the fraction of time spent in level n at ergodicity, apart from terms of order $O(\lambda^2/M^2)$, is given for $n \geq 1$ by

$$\frac{E\left[\int_0^D I_n(y) dy\right]}{D + M/\lambda} = \frac{\int_0^D r_n(y) dy}{D + M/\lambda}$$

From the familiar $\lim_{M \rightarrow \infty} (1 + x/M)^M = e^x$, one has that the p.g.f. of the number in the system at ergodicity is for fixed D ,

$$\begin{aligned} \pi(u, \infty) &= \lim_{M \rightarrow \infty} \left(\frac{M/\lambda + \int_0^D g(u, y) dy}{D + M/\lambda} \right. \\ &\quad \left. + O(\lambda^2/M^2) \right)^M \\ &= \exp\left[-\lambda \int_0^D \{1 - g(u, y)\} dy\right]. \end{aligned}$$

Since the duration of the holding times may be infinite, one obtains

$$\pi(u, \infty) = \exp\left[-\lambda \int_0^\infty \{1 - g(u, y)\} dy\right]. \tag{2}$$

A more formal demonstration of (2) can be obtained from a continuous infinite product representation of $\pi(u, \infty)$.

Note that for the ordinary $M/\bar{G}/\infty$ system with batch size $K = 1$ and service time c.d.f. $A_T(x)$, one has

$$g(u, y) = A_T(y) + u\bar{A}_T(y). \tag{3}$$

Hence

$$1 - g(u, y) = \bar{A}_T(y)(1 - u)$$

and one obtains from (2) the classical result $\pi(u, \infty) = \exp[-\lambda E\{T\}(1 - u)]$. (See for example Tijms [6].) Equation (2) implies at once that:

Theorem. For a process $N(t)$ with batch poisson arrivals of rate λ , independent identical random impulse response $g(u, t)$ for each batch and infinite number of servers, the ergodic distribution of the number of items in the system is a compound Poisson distribution with p.g.f. given by (2).

The basic result (2) can be extended by an identical argument to the continuum case for which

$$X(t) = \sum_{k=-\infty}^{k=+\infty} X_k^*(t - \tau_k). \tag{4}$$

Here, as above, the summands are i.i.d. stochastic response functions which go to zero as t goes to infinity, and (4) describes the superposition of such response functions to impulses at Poisson epochs. If $\phi(s, \infty) = E\{\exp\{-sX(\infty)\}\}$ one has

$$\phi(s, \infty) = \exp\left[-\lambda \int_0^\infty \{1 - \psi(s, y)\} dy\right]$$

where $\psi(s, y) = E\{\exp\{-sX_k^*(y)\}\}$. In the special case where $X_k^*(y)$ is deterministic and exponential this result coincides with Keilson and Mirman [3]. An extensive discussion of the general deterministic case may be found in Rice [4]. The result extends to multivariate processes of the form (4).

2. The ergo

From (2)
 $E\{N(\infty)\}$

If $N^*(t)$ is associated

$$E\{N(\infty)\}$$

In the same

$$\text{Var}\{N(\infty)\}$$

so that

$$\text{Var}\{N(\infty)\}$$

3. M^{batch}/λ

Until now about $N^*(t)$ independent helpful to special case served inde tially distrib size is exact

$$g(u, y) =$$

so that for

$$f_K(u) = \int_0^\infty$$

From the evaluate

$$h(K, r) =$$

from the in tion. It foll

$$\pi(u, \infty) =$$

This may

2. The ergodic mean and variance

From (2) one has at once

$$E[N(\infty)] = \lambda \int_0^\infty g_\nu(1, y) dy$$

$$= \lambda \int_0^\infty \sum_0^\infty nr_n(y) dy.$$

If $N^*(t)$ is the decreasing number in the system associated with each Poisson epoch, then

$$E[N(\infty)] = \lambda \int_0^\infty E[N^*(y)] dy. \tag{5}$$

In the same way, one finds that

$$\text{Var}[N(\infty)] = \lambda \int_0^\infty [g_{\nu\nu}(1, y) + g_\nu(1, y)] dy$$

so that

$$\text{Var}[N(\infty)] = \lambda \int_0^\infty E[N^{*2}(y)] dy. \tag{6}$$

3. $M^{\text{batch}}/M/\infty$

Until now no assumptions have been made about $N^*(t)$ other than that batches are served independently. As a check on the formalism, it is helpful to treat by an independent method the special case $M^{\text{batch}}/M/\infty$, where each customer is served independently and lifetimes are exponentially distributed. Let us also suppose that batch size is exactly K . Then, as for (3)

$$g(u, y) = [(1 - e^{-\theta y}) + u e^{-\theta y}]^K$$

so that for (2) one must evaluate

$$f_K(u) = \int_0^\infty [(1 - e^{-\theta y}) + u e^{-\theta y}]^K dy.$$

From the Binomial expansion, one must then evaluate

$$h(K, r) = \int_0^\infty (1 - e^{-\theta y})^{K-r} e^{-r\theta y} dy$$

$$= [\theta r^K C_r]^{-1}$$

from the integral representation of the Beta function. It follows that

$$\pi(u, \infty) = \exp\left[-\frac{\lambda}{\theta} \sum_1^K \frac{1}{r} + \frac{\lambda}{\theta} \sum_1^K \frac{u^r}{r}\right]. \tag{7}$$

This may be seen to coincide with the p.g.f. ob-

tained for $M^b/M/\infty$ by analysis via a birth-death process for general batch size distribution. Let $a_n = P[K = n]$ and $p_n(t) = P[N(t) = n]$. One then has from the forward Kolmogorov equations

$$\frac{d}{dt} p_n(t) = -(\lambda + n\mu) p_n(t) + \lambda (p_n(t))^* (a_n) + (n+1)\mu p_{n+1}(t).$$

The use of generating functions then gives

$$\frac{\partial}{\partial t} \pi(u, t) = -\lambda [1 - \alpha(u)] \pi(u, t) + \mu(1 - u) \frac{\partial}{\partial u} \pi(u, t),$$

i.e.

$$(1 - u) \frac{\partial}{\partial u} \log \pi(u) = \frac{\lambda}{\mu} [1 - \alpha(u)].$$

One has finally

$$\pi(u) = \exp\left[\frac{\lambda}{\mu} \int_1^u \frac{[1 - \alpha(w)]}{1 - w} dw\right]. \tag{8}$$

This may be seen to coincide with (7) when $\alpha(u) = u^K$.

4. $M^A/G/\infty$

For constant batch size and general holding times T with c.d.f. $A_T(x)$, one has

$$g(u, y) = [A_T(y) + u\bar{A}_T(y)]^A$$

so that

$$\pi(u, \infty) = \exp\left[\lambda \int_0^\infty [g(u, y) - g(1, y)] dy\right]$$

$$= \exp\left[\lambda \int_0^\infty \{[A(y) + u\bar{A}(y)]^A - [A(y) + \bar{A}(y)]^A\} dy\right].$$

Hence

$$\pi(u, \infty) = \exp\{-\theta [1 - \beta(u)]\} \tag{9}$$

where

$$\theta = \lambda E[\max(T_1, T_2, \dots, T_A)]$$

$$= \lambda \int_0^\infty [1 - A^A(y)] dy. \tag{10}$$

If we use the binomial expansion and denote the binomial coefficients by C_k^K , we obtain

$$\beta(u) = \frac{\sum_1^K u^k C_k^K \int_0^\infty \bar{A}_T^k(y) A_T^{K-k}(y) dy}{\int_0^\infty [1 - A_T^K(y)] dy} \quad (11)$$

Note that only when $K = 1$ is the distribution of $N(\infty)$ independent of the lifetime distribution. Note also that for constant batch size K one has from differentiation

$$E[N(\infty)] = \lambda K E[T], \quad (12)$$

$$\text{Var}[N(\infty)] = \lambda K E[T] + \lambda K(K-1) \times \int_0^\infty [1 - A_T(y)]^2 dy, \quad (13)$$

in agreement with Tijms (1986). The numerical evaluation of the distribution of $N(\infty)$ from (9) can be obtained algorithmically from

$$\exp\{-\theta[1 - \beta(u)]\} = e^{-\theta} \sum \frac{\theta^n}{n!} \beta^n(u)$$

and n -fold convolution of the lattice distribution for $\beta(u)$ with itself. It is clear from the compound Poisson character of (9) that for any holding time distribution with finite mean, when K is fixed and λ becomes large, the distribution of $N(\infty)$ be-

comes normal. Normality does not set in with increasing batch size alone since $\pi(u, \infty)$ is not the K -th power of a p.g.f. The normality will be present when $\theta = \lambda E[\max(T_1, T_2, \dots, T_k)]$ is large.

Acknowledgement

This study was initiated at and supported in substantial part by the Wm. E. Simon School of the University of Rochester. The authors are indebted to S. Graves, L. Servi, U. Simuta and D. Nakazato for helpful comments.

References

- [1] M. Brown and S.M. Ross, "Some results for infinite server Poisson Queues" *J. Appl. Prob.* 6, 604-611 (1969).
- [2] M.L. Chaudry and J.G.C. Templeton, *A First Course in Bulk Queues*, Wiley, New York.
- [3] J. Keilson and N.D. Mirman, "The second order distribution of integrated shot noise", *IRE Transactions of Information Theory*, 1959, 75-77.
- [4] S.O. Rice, "Mathematical analysis of random noise", *Bell System Technical Journal* 23 (1945).
- [5] D.N. Shanbhag, "On infinite server queues with batch arrivals", *J. Applied Prob.* 2, 974-979 (1966).
- [6] Henk C. Tijms, *Stochastic Modelling and Analysis*, Wiley, New York, 1986.

A DIS

J. KEILSON

GTE LABORATORY

L.D. SIVAN

GTE LABORATORY

Received

Revised

For manuscript

the preparation

of this paper

and the preparation

of this paper

and the preparation

of this paper

1. Introduction

Over the past few years, the importance of queueing theory has increased. Ramalhoto (1986) equates the term. For many applications, the Law is applied to distribution setting requirements. The distribution observed by Mann and Zipkin [17], its importance set forth.

Suppose that in a queueing system, there is a Poisson arrival process with number of arrivals N and let T be the service time of a customer in a queueing system. The p.g.f. of N is

* 40 Sivan